

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FOOD CONTACT SURFACE SANITIZATION DIRECTIONS

Sanitization: Recommended for use in restaurants, dairies and bars. When used as directed this product is an effective sanitizer at an active quaternary concentration of 200 - 400 ppm in public eating establishments and dairies against *Escherichia coli* (ATCC 11229) *Escherichia coli* O157:H7, *Staphylococcus aureus*, *Yersinia enterocolitica*, *Listeria monocytogenes*, *Salmonella typhi*, *Shigella sonnei*, *Vibrio cholera*, Methicillin resistant *Staphylococcus aureus* (MRSA), Vancomycin resistant *Enterococcus faecalis* (VRE), *Campylobacter jejuni* (ATCC 29428) and *Klebsiella pneumoniae*. This product is an effective sanitizer against *Klebsiella pneumoniae* at a 200 ppm active quaternary concentration when diluted in water up to 500 ppm hardness (CaCO₃). At an active quaternary concentration of 300 ppm, it is an effective sanitizer against *Klebsiella pneumoniae* when diluted in water up to 650 ppm hardness (CaCO₃). Remove all gross food particles and soil from areas, which are to be sanitized, with a good detergent, pre-flush, pre-soak or pre-scrape treatment. Rinse with a potable water rinse. Prepare a fresh solution daily or more frequently as soil is apparent.

To sanitize pre-cleaned and potable water-rinsed, non-porous public eating establishment and dairy food contact surfaces: prepare a 200 - 400 ppm active quaternary solution by adding 1 - 2 ounces of **Sani-Guard** to 4 gallons of water, 2.5 - 5 ounces to 10 gallons of water, or 5 - 10 ounces to 20 gallons of water.

To sanitize pre-cleaned mobile items in public eating establishments (drinking glasses, dishes, eating utensils) immerse in a 200 - 400 ppm active quaternary solution for at least 60 seconds making sure to immerse completely. Remove items, drain the use-solution from the surface and air dry. Do not rinse.

To sanitize pre-cleaned public eating establishment surfaces (counters, tables, finished wood or plastic cutting boards), apply a 200 - 400 ppm active quaternary solution with a cloth, sponge, low pressure coarse sprayer or hand pump trigger sprayer making sure that the surface remains completely wet for at least 60 seconds and let air dry. Do not rinse.

To sanitize pre-cleaned dairy processing equipment (tanks, piping, pasteurizers, milking equipment) apply a 200 - 400 ppm active quaternary solution with a low pressure coarse sprayer or circulate a 200 - 400 ppm active quaternary solution throughout the equipment. Surfaces must remain thoroughly wet for at least 60 seconds, followed by adequate draining and air drying. Do not rinse. At 1 ounce per 4 gallons, this sanitizer fulfills the criteria of Appendix F of the Grade A Pasteurized Milk Ordinance 1978 Recommendations of the U.S. Public Health Services in waters up to 650 ppm of hardness calculated as CaCO₃ when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*.



Sani-Guard™ SANITIZER

ACTIVE INGREDIENTS	
Alkyl (60% C14, 30% C16, 5% C12, 5% C18)	
dimethyl benzyl ammonium chlorides.....	5.0%
Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chlorides.....	5.0%
INERT INGREDIENTS.....	90.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN DANGER

FIRST AID - Have the product container or label with you when calling a poison control center or doctor, or going for treatment.
If in eyes - Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed - Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If inhaled - Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
NOTE TO PHYSICIAN - Probable mucosal damage may contraindicate the use of gastric lavage.

See side panel for additional precautionary statements.

EPA REG. NO. 1839-86-34370 BATCH:
EPA EST. NO. 34370-LA-001 NET CONTENTS: 1 U.S. GALLON

WISCONSIN STATE DIVISION OF HEALTH
DIRECTIONS FOR EATING ESTABLISHMENTS
1. Scrape and pre-wash utensils and glasses whenever possible.
2. Wash with a good detergent or compatible cleaner.
3. Rinse with clean water.
4. Sanitize in a solution of 1 ounce to 4 gallons of water (200 ppm active quaternary). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drain board to air dry.
6. Fresh sanitizing solution should be prepared at least daily or when visibly soiled or diluted.
NOTE: A clean potable water rinse following sanitization is not permitted under Section HFS 196.13 of the Wisconsin Administrative Code.
Sanitization of beverage dispensing equipment:
1. Wash equipment with a compatible detergent and rinse with potable water prior to sanitizing.
2. Fill equipment with a use-solution of 0.25 - 0.5 ounces of **Sani-Guard** per gallon of water (or equivalent dilution).
3. Allow use-solution to remain in equipment for at least 60 seconds.
4. Drain thoroughly and allow to air dry before reuse. Do not rinse with potable water.

(Not for Use in California) Entryway sanitizing systems: To prevent cross contamination from area to area, set the system to deliver 0.51 to 1.02 ounces per gallon of water (or equivalent use dilution) (400 to 800 ppm active quaternary) of sanitizing solution. The spray should cover the entire path of the doorway. Set the system so that a continuous wet blanket of sanitizer solution is delivered to the floor. Do not mix other additives to the sanitizing solution.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

KEEP OUT OF REACH OF CHILDREN. **CORROSIVE.** Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. Wear goggles or face shield, rubber gloves, and protective clothing. Harmful if swallowed. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL

PESTICIDE STORAGE - Store in a dry place no lower in temperature than 50°F or higher than 120°F. Store in a tightly closed container in an area inaccessible to children.

PESTICIDE DISPOSAL - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL - Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container promptly after emptying.

Triple rinse as follows: Fill container 1/4 full with water and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times. Then offer for recycling or reconditioning. If not available, puncture and dispose in a sanitary landfill.

SPILL OR LEAK PROCEDURES

Small spills may be mopped up, flushed away with water or absorbed on some absorbent material and incinerated. Large spills should be contained, the material then moved into containers and disposed of by approved methods for hazardous wastes.

Sold by: **WECHEM, INC.**
5734 Susitna Drive
Harahan, LA 70123
(504) 733-1152
800-426-0512
FAX: (504) 733-2218
www.wechem.com

7 38435 20724 3